

AD _____

COOPERATIVE NUMBER DAMD17-93-V-3002

TITLE: Research in Experimental Antimalarial Chemotherapy

PRINCIPAL INVESTIGATOR: Dr. Ayoade M. J. Oduola

CONTRACTING ORGANIZATION: Walter Reed Army Institute of Research
Washington, DC 20307-5100

REPORT DATE: January 1995

TYPE OF REPORT: Final

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19970121 143

DTIC QUALITY INSPECTED 1

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE January 1995	3. REPORT TYPE AND DATES COVERED Final (1 Oct 92 - 30 Dec 94)		
4. TITLE AND SUBTITLE Research in Experimental Antimalarial Chemotherapy		5. FUNDING NUMBERS DAMD17-93-V-3002		
6. AUTHOR(S) Dr. Ayode M. J. Oduola				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research Washington, DC 20307-5100		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, MD 21702-5012		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) The primary objectives of the project on Research on Antimalarial Chemotherapy: "Novel approach to combination therapy in multidrug resistant malaria" focused on confirmation of reversal of chloroquine resistance in isolates of <i>P. falciparum</i> from Nigeria and molecular biology, pharmacokinetics and biochemistry of the reversal phenomenon. The major objectives of the project were accomplished. Efforts during the project resulted in modification of standard <i>in vitro</i> techniques for continuous cultivation of <i>P. falciparum</i> in Nigeria. The modified technique is now in routine use at the cooperative malaria laboratory in the department of Pharmacology and Therapeutics University of Ibadan, Nigeria. Reversal of mefloquine resistance with penfluridol and reversal of chloroquine resistance with promethazine (WRBL 50601), chlorpheniramine, diphenhydramine and pyrilamine were identified and published in 1991 and 1992. The findings on reversal of chloroquine resistance with promethazine and chlorpheniramine represent the first demonstration of the reversal phenomenon with drugs prescribed as adjunct in chemotherapy of malaria in an endemic area. Subsequent efforts were allocated to studies on pharmacokinetics and clinical efficacy of potential combinations of chlorpheniramine with antimalarial drugs in human volunteers and animal models. Other studies accomplished during the period include clinical efficacy of artemether, mefloquine and suphadoxine-pyrimethamine. Nine manuscripts have been published, two are being prepared for publication.				
14. SUBJECT TERMS Chemotherapy		15. NUMBER OF PAGES 21		16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

✓ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

____ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

____ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

✓ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.


 02/14/96
PI - Signature Date

Table of Contents

	Page
I. Introduction and Summary of Achievement of the Objectives of the Grant	1
a. Objectives of the project	1
b. Achievement of the Grant	1
II. Research Activities	2
a. Studies on the pharmacokinetics of quinine combinations	2
b. Analysis	3
c. Studies on the effect of chlorpheniramine on <i>in vitro/in vivo</i> susceptibility of <i>Plasmodium falciparum</i> to chloroquine in south west Nigeria	4
III. Publications	6
IV. Staff Receiving Remuneration and Development Training	8
V. In-House Research Training	8

I Introduction and Summary of Achievement of the Objectives of the Grant

Ia. The objectives of the project are:

1. To evaluate the clinical efficacy of combination therapy with chloroquine and reversing compounds (chlorpheniramine or promethazine) in treatment of chloroquine resistant malaria.
2. To evaluate and confirm reversal of resistance to chloroquine, quinine, quinidine and mefloquine with verapamil or penfluridol in isolate and clones of *P. falciparum* obtained from Nigeria. Correlate reversal of CQR with clinical response of the patients and use the unique reversal with verapamil in enhancing a technique for detection of CQR *P. falciparum*.
3. To evaluate effects of co-administration of reversing compounds on pharmacokinetic properties of standard antimalarial drug.

Ib. Achievement of the Grant

This represents the final report on project (Cooperative Agreement DAMD17-93-V-3002) WHO/TDR ID # 890178 (Oct. 1992 to Dec. 1994). In order to avoid repetition of work already enumerated in publications and earlier reports, a brief summary of the achievements of the project is provided with emphasis on achievements not previously reported. The primary objectives of the project on Research on Antimalarial Chemotherapy: "Novel approach to combination therapy in multidrug resistant malaria" remained unchanged. It focused on confirmation of reversal of chloroquine resistance in isolates of *P. falciparum* from Nigeria and studies on molecular biology, pharmacokinetics and biochemistry of the reversal phenomenon. The major objectives of the project were accomplished during the grant period.

Efforts during the project resulted in modification of standard *in vitro* techniques for continuous cultivation of *P. falciparum* in Nigeria. The modified technique is now in routine use at the cooperative malaria laboratory in the department of Pharmacology and Therapeutics University of Ibadan, Nigeria. Reversal of mefloquine resistance with penfluridol and reversal of chloroquine resistance with promethazine (WRBL 50601), chlorpheniramine, diphenhydramine and pyrilamine were identified and constitute part of reports published in 1991 and 1992. Subsequently, reversal of chloroquine resistance with promethazine and chlorpheniramine have been confirmed in additional isolates of *P. falciparum* obtained from patients in Nigeria. This finding represents the first demonstration of the reversal phenomenon with drugs prescribed as adjunct in chemotherapy of malaria in an endemic area. The significance of these observations influenced allocation of subsequent efforts to studies on pharmacokinetics and clinical efficacy of potential combinations of chlorpheniramine with antimalarial drugs in human volunteers and animal models. Chlorpheniramine was selected for further studies because of the milder side effects compared with promethazine. Major activities not previously reported or published are outlined below.

- 1. Confirmation of reversal of chloroquine resistance and identification of reversal of quinine resistance with chlorpheniramine and promethazine in isolates of *P. falciparum* adapted to continuous culture in Nigeria.
- 2. Studies on Pharmacokinetics interactions of chlorpheniramine and acetaminophen with quinine.
- 3. Studies on Pharmacokinetics interaction of promethazine with chloroquine.
- 4. Studies on the clinical efficacy of a combination of chloroquine with chlorpheniramine in the treatment of CQR malaria.

• In addition to these studies on the clinical application of the reversal phenomenon, efforts were devoted to studies on.

- 1. Clinical studies on efficacy of artemether, mefloquine and suphadoxine-pyrimethanine.
- 2. Studies on parasite ring viability following antimalarial drug administration.
- A total of nine manuscripts have been published while additional two are being prepared for publication. A total of eleven students (7 Ph.D and 5 M.Sc) were trained during the period of the grant. In addition 8 technical staff received training in various techniques for studies on malaria. One WHO post doctoral fellow (Dr. I. Obarisiagbon) started training on techniques for continuous cultivation of *P. falciparum* and isoenzyme analysis for studies on molecular biology of chloroquine resistant *P. falciparum*.

II Research Activities

STUDIES ON THE PHARMACOKINETICS OF QUININE COMBINATIONS.

- The Pharmacokinetic disposition of quinine and combinations with either tetracycline or paracetamol were studied after oral administration to healthy African subjects. All study protocols were approved by the Joint University College Hospital and College of Medicine University of Ibadan ethical review committee. Twenty-one healthy male adults aged between 21 and 39 years volunteered and participated in the study after giving written informed consents. The volunteers were non-smokers who consumed alcohol only occasionally. They weighed between 58 and 74kg. Each subject was selected for participation based on normal findings on physical examination and normal urinalysis, biochemical and hematological tests. In addition, they did not take any medication for 4 weeks prior to participation and during the period of the study.
- The subjects fasted overnight before the study up till 3 hours after drug administration. A fore-arm vein was cannulated for blood sampling and kept patent with heparinized normal saline (heparin 5iu/ml). The Drug(s) were administered at 3 week intervals by a randomized cross-over design.
- Drug regimen:
- (i) Quinine/Paracetamol

- Six subjects received either oral quinine sulphate (600mg) or oral paracetamol (1000mg) 6 hourly for the first 3 doses followed by a combination of paracetamol (1000mg) and quinine sulphate (600mg) after a 12 hour (overnight) fast.
- This regimen was designed to simulate as closely as possible the usual self treatment history of the typical patient before seeking medical treatment.
- (ii) Quinine/Chlorpheniramine
- Eight subjects received either oral quinine sulphate (600mg) or oral chlorpheniramine (4mg) followed by a combination of quinine sulphate (600mg) and chlorpheniramine (4mg).
- (iii) Quinine/Tetracycline
- Seven subjects received either oral quinine sulphate (600mg) or tetracycline (250mg) followed by a combination of quinine sulphate (600mg) and tetracycline (250mg).
- Sample collection.
- Venous blood was obtained before and at 0.25, 0.5, 0.75, 1.0, 2.0, 3.0, 4.0, 6.0, 8.0, 10.0, 12.0, 24.0, 36.0 and 48.0 hours after drug administration.
- The plasma and red cells were separated after centrifuging at 1200 rpm for 10 minutes and stored at -20°C . Each subjects blood pressure and pulse rate were monitored every hour for the first 12 hours after drug administration and subsequently at the time of blood sampling. The subjects were questioned for any subjective feeling during the first 12 hours after drug administration.

Iib. Analysis:

- The concentration of quinine in plasma was measured by a high-performance liquid chromatography method using a reverse phase column and a fluorescence detector (Sowunmi and Salako 1992).
- The intra and inter assay co-efficients of variation were 5.5 and 2.5% for the concentrations of 10ng/ml and 10ug/ml respectively. Recoveries over the concentration range of 10ng/ml to 10ug/ml were 93-100%. The lower limit of detection was 20ng/ml. Calibration plots were linear ($r = 0.9897$) up to 8 ug/ml.
- The pharmacokinetic parameters calculated from the concentration time data are presented in tables. The pharmacokinetic parameters of quinine (500mg base) following a single intramuscular injection (7 volunteers) or oral (11 volunteers) administration in healthy Nigerian volunteers are shown on Table 1.
- There were no significant differences ($p > 0.05$) in the maximum plasma concentration when quinine was administered alone or in combination with paracetamol ($C_{\text{max}} = 2.10 \pm 0.5 \text{ ug/ml}$ vs $2.50 \pm 0.8 \text{ ug/ml}$; $T_{\text{max}} = 4.3 \pm 2.8 \text{ h}$ vs $4.4 \pm 2.2 \text{ hr}$ respectively). The mean elimination half life of quinine when administered alone ($10.5 \pm 3.7 \text{ h}$) was similar to that obtained when co-administered with paracetamol ($9.5 \pm 3.7 \text{ h}$). The oral clearance of quinine and the area under the plasma concentration time curve was also not significantly

different when quinine was administered alone 0.26 ± 0.09 L/h/kg and 39.1 ± 15.1 ug/ml.h respectively) or when co-administered with paracetamol (0.23 ± 0.03 L/h/kg and 40.7 ± 6.1 ug/ml.h respectively) (Table 2). The data obtained from the subjects studied (n=6) indicate that co-administration of paracetamol with quinine in therapeutic doses does not significantly affect the disposition kinetics of quinine in healthy young adults.

- The interaction between quinine and the resistance reversing compound chlorpheniramine was studied in 8 subjects. Plasma quinine C_{max} was increased from 2.9 ± 0.7 mg/l when quinine was administered alone to 3.1 ± 1.1 mg/l after co-administration of chlorpheniramine. Similarly the elimination half life increased from 12.6 ± 5.5 h. to 13.5 ± 4.6 h when chlorpheniramine was co-administered with quinine. The volume of distribution was however reduced from 2.63 ± 0.85 L/kg to 2.39 ± 0.55 L/kg following administration of quinine / chlorpheniramine combination. These differences were not statistically significant. There were however significant reductions in quinine AUC₀₋₄ and plasma clearance (Cl_p) by the combination. (Table 3).
- Combination of quinine/chlorpheniramine significantly increased quinine profile in the red cell. There were significant increases in C_{max}, T_{1/2}, T_{max} and AUC values following co-administration of chlorpheniramine (Table 4). The combination was however well tolerated in all subjects studied.
- Co-administration of tetracycline with quinine did not produce any significant changes in quinine pharmacokinetic profile. (Table 5).
- Data obtained from studies on chloroquine/promethazine interaction are currently being analysed.

IIc. STUDIES ON THE EFFECT OF CHLORPHENIRAMINE ON IN-VITRO/IN-VIVO SUSCEPTIBILITY OF PLASMODIUM FALCIPARUM TO CHLOROQUINE IN SOUTH WEST NIGERIA.

IN VIVO STUDY.

- Patients (under 5 years of age) with clinical symptoms and microscopically confirmed *P. falciparum* infection were randomly allocated into one of two treatment groups if following criteria were fulfilled.
 - (1) Pure asexual *P. falciparum* infection > 8,000 parasites per ul blood.
 - (2) Negative history of antimalarial drug ingestion in preceding 3 weeks.
 - (3) Negative dill glazko and lignin tests.
 - (4) Parent/guardian consent.
- One group received chloroquine alone (25mg/kg body weight) given in 3 doses, while the other group received a combination of chloroquine (given as above) and chlorpheniramine (6mg in three divided doses). Parasitological and clinical response of

each patient to treatment were monitored using the WHO extended 28-day test. Thick and thin blood films were prepared for each patient daily for 3 consecutive days and subsequently weekly till day 28. Blood samples were obtained from each patient before treatment and used for WHO standard schizont inhibition assay.

***IN-VIVO* RESPONSE**

Nineteen (19) out of the initial 30 patients enrolled completed the study. Eight (8) patients were treated with chloroquine alone while eleven (11) patients received the combination of chloroquine and chlorpheniramine. Cure rate by day 28 was 38% and 64% in the chloroquine and chloroquine/chlorpheniramine group respectively.

***IN-VITRO* STUDY**

Blood samples were obtained from each patient before treatment. An aliquot of each sample was used for *in vitro* drug susceptibility testing using a modification of the WHO schizont inhibitory assay (Reckmann et al., 1978). Ninety-six well microtitre plates containing serial dilutions of chloroquine were modified to incorporate fixed concentrations of chlorpheniramine ($1 \times 10^{-6} \text{M}$) or verapamil ($1 \times 10^{-6} \text{M}$). Each assay was monitored for 24hr-36hrs. At the end of each experiment giemsa-stained thick blood films were prepared and examined microscopically.

Sixteen (16) of the schizont inhibition assays were successful. Two (2) of the isolates were sensitive to chloroquine *in vitro* using a cut off concentration of 3.55 ug/l. The minimum inhibitory concentrations for chloroquine against the isolates was 2.10 ng/ml. Thirteen (13) isolates were resistant to chloroquine by this criteria. The minimum inhibitory concentrations for chloroquine against the isolates ranged from 6.20 ng/ml to 166.66 ng/ml. Simultaneous exposure of resistant parasites to a combination of chloroquine and a reversing compound increased susceptibility of 7 of the isolates to chloroquine. In contrast, combination of chloroquine with the reversing compound produced an increase in MIC in 2 sensitive isolates (Table 6). Although the implication of a decrease in the activity of chloroquine against a sensitive isolate in the presence of a reversing compound is not clear, it has been previously observed in cultured isolates (Kyle et al., 1990)

Complete *in vivo* and *in vitro* data were obtained for 10 patients. Six (6) of them received chloroquine alone while 4 patients were treated with the combination of chloroquine and chlorpheniramine (CQ/CP). In the CQ group 3 patients were cured of infection while recrudescence occurred in 2 patients and infection in one patient failed to respond to treatment. *In vitro* data for these patients showed that 3 isolates were sensitive and 3 isolates were resistant to chloroquine. Chlorpheniramine did not exhibit any effect on the activity of chloroquine against these resistant parasites *in vitro*. In the CQ/CP group (4), three (3) of the patients were cured of infection while infection in the fourth patient failed to respond to therapy. *In vitro* data for the patients showed that 3 of the isolates of *P. falciparum* were resistant to chloroquine while only one isolate was sensitive.

Chlorpheniramine markedly reduced MIC for chloroquine against the resistant parasites *in vitro*.

The results indicate that chlorpheniramine enhances the intrinsic antimalarial activity of chloroquine against chloroquine resistant parasites. However a larger study group is required to confirm this observation against chloroquine resistant malaria infections. The study is currently being repeated in a larger population using a double blind placebo controlled design. Results of the study would constitute an addendum to this report.

III PUBLICATIONS

- (1) ODUOLA, A.M.J, Sowunmi, A., Milhous, W.K., Kyle D.E., Martin R.K. Walker O and Salako L.A. (1992) Innate Resistance to New Antimalarial Drugs in *Plasmodium falciparum* from Nigeria. Transaction of the Royal Society of Tropical Medicine and Hygiene. 86:123-126.
- (2) Sowunmi, A., ODUOLA, A.M.j., Salako, L.A., Ogundahunsi, O.A.T. and Laoye, O.J. (1992). The relationship between response of *Plasmodium falciparum* to mefloquine in African children and its sensitivity *in vitro*. Transaction of the Royal Society of Tropical Medicine and Hygiene. 86: 368-371.
- (3) ODUOLA, A.M.J., Omitowoju, G.O., Garena, L., Kyle, D.E., Milhus, W.k., Sowunmi, A. and Salako, L.A. (1993). Reversal of mefloquine resistance with penfluridol in isolates of *Plasmodium falciparum* from southwest Nigeria. Transaction of the Royal Society for tropical medicine and Hygiene 87:81-83.
- (4) ODUOLA, A.M.J., Ogundahunsi, O.A.T. and Salako, L.A. (1992). Continuous cultivation and drug susceptibility testing of *Plasmodium falciparum* in malaria endemic area. Journal of Protozoology 39(5) 605-608.
- (5) A. Sowunmi, L.A. Salako, A.M.J. ODUOLA, O, Walker, J.A. Akindele and O.A.T. Ogundahunsi. (1993). Neuropsychiatric side effects to mefloquine in Africans Transactions of the Royal society for tropical medicine and Hygeine 87:462-463.
- (6) A. Sowunmi, J.A. Akindele, G.O. Omitowoju, A. Omigbodum, ODUOLA A.M.J and Salako L.A (1993). Intramuscular sulphadoxine- Pyrimethamine in uncomplicated chloroquine-resistant falciparum malaria during pregnancy. Transactions of the Royal Society for Tropical Medicine and hygiene 87:472.
- (7) Sowunmi, A., ODUOLA, A.M.J., Ilesanmi, A.O. and Salako, L.A. 1994. Open comparison of artemether and mefloquine in uncomplicated *Plasmodium falciparum* hyperparasitaemia in children. Annal Trop. Paediatr. Accepted for Publication.

- (8) Salako, L.A., Walker, O., Sowunmi, A., Omokhodion S. I., Adio. R.A. and **ODUOLA**, A.M.J. 1994. Artemether in severe and cerebral malaria in Nigerian Children. Trans. Roy. Soc. Trop. Med. Hyg., 88: Supplement 1. 13-15
- (9) Sowunmi, A. and **ODUOLA**, A.M.J. 1994. Open comparison of mefloquine, mefloquine-sulfadoxine-pyrimethamine and chloroquine in acute uncomplicated falciparum malaria in children. Trans. Roy. Soc. Trop. Med. Hyg., 89:303.

IV Staff Receiving Remuneration and Development Training

- (i) **Mrs. R.A. Adio** (Chief technologist) received training between July-and August-1991 on application of the isolated perfused rat liver system. Her training was at the department of Pharmacology, University of Sheffield and Division of Experimental Therapeutics, WRAIR Washington D.C. USA.
- (ii) **Mr. K.T. Babalola** (AIMLT, M.Sc). Completed training in the techniques for preparation of DNA from isolates of *P. falciparum*.
- (iii) Three technicians with the national diploma in laboratory technology were trained in the laboratory (i) **Mr. A.O. Sijuade** was trained on the use of HPLC for analysis of antimalarial drugs in body fluids. (ii) **Ms. Oyelakin** was trained on techniques for continuous cultivation of *Plasmodium falciparum* and use of the WHO micro test kit. (iii) **Ms. Olaniran** was trained in techniques for continuous cultivation and drug susceptibility testing of *P. falciparum*.
- (iv) **Mr. Wole Oyebamiji**, a laboratory trainee assistant has been trained to assist in pharmacokinetics studies in human volunteers and routine microscopy.
- (v) **Mr. Aina (BSc) and Mr. Thomas (MSc)** were trained on techniques for enzyme determination and DNA extraction respectively. In addition they were both trained on techniques for continuous cultivation and drug susceptibility testing of *P. falciparum*.
- (vi) **Miss Sanni** a trainee laboratory technologist is being trained on the technique for continuous cultivation of *Plasmodium falciparum* and use of the WHO micro test kit.

V. In-House Research Training

A total of 11 students (7 Ph.D and 5 M.Sc) were trained or are being trained.

- 1 **Mr. B. Umotong (Ph.D)**: Primarily attached to the Department of Immunology received training in the use of phase contrast microscopy for IFA. He is currently in the final stages of defending a thesis on "Immunological studies of Nigerians infected with chloroquine resistant *P. falciparum*".
- 2. **Mr. Hedo (Ph.D)**: Primarily attached to the Department of Immunology received training in the use of phase contrast microscopy for IFA. He is currently in the final stages of defending a thesis on. "Immunological parameters and severity of sickle cell anemia in Nigerian adults".

3. **Mr. Bola Taiwo (Ph.D):** Received training on technique for evaluating cardiotoxicity of reversing compounds alone and in combination with chloroquine using experimental animal models.
4. **Mr. O.A.T. Ogundahunsi (Ph.D)** Received additional training in techniques for (a) *in vitro* continuous cultivation and drug susceptibility testing of *P. falciparum* (b) screening potential resistance reversing compounds (c) biological assay for evaluating reversing ability of drugs after oral absorption and potential biotransformation (d) Use and modification of the WHO micro test Kit for rapid diagnosis of CQ resistant malaria.
5. **Mr. V. Nwagbarocha (M.Sc):** Completed studies culminating in the award of M.Sc (Pharmacology & Therapeutics). His thesis supported by the lab study project is titled "*In vitro* cultivation of *P. falciparum* using plasma free culture medium".
6. **Dr. C.O Kawesha (M.Sc)** A WHO/TDR trainee received training on conduct of clinical efficacy studies and Pharmacokinetics. His thesis on "Pharmacokinetic interactions of quinine and tetracycline" was successfully defended for the degree of M.Sc in Clinical Pharmacology.
7. **Mr. E Achidi (Ph.D)** Primarily attached to the Department of immunology received training in the use of phase microscopy for IFA.
8. **Miss. G.O. Omitowoju (Ph.D)** Obtained a M.Sc (Pharmacology & Therapeutics) in 1991 based on studies conducted in the lab. She is currently registered for a doctorate with the group.
9. **Mrs. Charles-Davies (Ph.D)** primarily attached to the Department of Chemical pathology. She received training on the use of phase contrast microscopy.
10. **Mr. Wale Oduola** is currently registered for a M.Sc degree in Pharmacology.
11. **Mr. Aina** is registered for an M.Sc degree in Pharmacology based on studies to be conducted in the Laboratory.

TABLE 1

PHARMACOKINETIC PARAMETERS OF QUININE (500 MG BASE) AFTER A SINGLE INTRAMUSCULAR INJECTION OR ORAL ADMINISTRATION IN HEALTHY NIGERIAN ADULTS.

Parameter	Intramuscular Quinine (n = 7)	Oral Quinine (n = 11)
C_{max} (mg/l)	4.50 + 1.30	2.90 + 0.50
t_{max} (hr)	1.70 + 1.10	3.30 + 0.80
$T_{1/2}$ (hr)	10.70 + 3.50	11.70 + 2.90
V_z (l/kg)	2.18 + 0.99	2.50 + 0.70
Cl_p (l/hr/kg)	0.15 + 0.09	0.15 + 0.04
$AUC_{0-\infty}$ (mg hr/l)	71.80 + 29.40	54.90 + 19.

TABLE 2a PLASMA PHARMACOKINETIC PARAMETERS FOR QUININE IN
6 HEALTHY SUBJECTS FOLLOWING ORAL ADMINISTRATION
OF 600 MG QUININE SULPHATE.

Volunteer #	Cmax (ug/ml)	Tmax (h)	t _{1/2} (h)	Vd (L/kg)	Clp (Oral) (L/h/kg)	AUC (ug/ml.h)
1	3.1	4	12.6	2.1	0.17	57.6
2	1.9	3	11.9	5.5	0.28	32.9
3	1.9	3	14.9	4.2	0.16	58.5
4	2.0	10	10.9	1.3	0.31	27.9
5	2.3	3	6.8	2.3	0.28	34.3
6	1.6	3	5.4	2.7	0.40	23.7
Mean	2.1	4.3	10.5	3.0	0.26	39.1
s.d.	0.5	2.8	3.7	1.5	0.09	15.1

TABLE 2b PLASMA PHARMACOKINETIC PARAMETERS FOR QUININE IN 6 HEALTHY
SUBJECTS FOLLOWING ORAL ADMINISTRATION OF 600 MG QUININE SULPHATE
AND 1000 MG PARACETAMOL.

Volunteer #	Cmax (ug/ml)	Tmax (h)	t _{1/2} (h)	Vd (L/kg)	Clp (Oral) (L/h/kg)	AUC (ug/ml.h)
1	3.5	1.5	8.9	2.2	0.20	48.4
2	1.5	6	8.8	2.4	0.25	37.0
3	3.4	4	12.1	1.8	0.24	39.2
4	1.8	8	13.7	2.1	0.18	48.0
5	2.5	3	7.0	2.1	0.25	38.3
6	2.3	4	6.9	1.8	0.28	33.5
Mean	2.5	4.4	9.5	2.1	0.23	40.7
s.d.	0.8	2.2	2.7	0.2	0.03	6.1

TABLE 3a. PLASMA PHARMACOKINETIC PARAMETERS AFTER A SINGLE ORAL DOSE OF 500MG QUININE BASE WITH 4MG CHLORPHENIRAMINE.

Subject	Weight kg	Cmax mg L-l	tmax h	t1/2 h	Vz Lkg-l	CLP Lhkg-l	AUC ₀ mgh L-l	AUC ⁰⁰⁰ Mg h L-l
1	67	2.9	4	18.0	2.38	0.09	6.68	79.8
2	62	3.1	3	9.2	2.39	0.18	42.8	79.8
3	60	2.2	4	15.8	3.10	0.13	53.1	59.9
4	74	1.9	6	9.4	3.31	0.24	25.8	27.1
5	61	4.4	4	19.8	1.71	0.06	111.2	134.1
6	64	3.0	1	8.4	2.25	0.18	38.8	41.2
7	58	4.4	6	10.4	1.87	0.11	72.6	75.6
8	62	3.4	3	17.3	2.16	0.08	81.5	69.1
mean	63.5	3.1	3.8	13.5	2.39	0.13	61.6	69.1
SD	5.0	1.1	1.6	4.6	0.55	0.06	27.3	34.0

TABLE 3b. PLASMA PHARMACOKINETIC PARAMETERS AFTER A SINGLE ORAL DOSE OF 5000MG QUININE BASE ALONE.

Subject	Weight kg	Cmax mg L-1	tmax h	t _{1/2} h	Vz Lkg ⁻¹	GLP Lhkg ⁻¹	AUC ₀ mgh L-1	AUC [∞] Mg h L-1
1	67	2.8	2	7.8	2.2	0.18	49.1	40.7
2	62	3.0	2	10.2	2.39	0.16	47.4	48.9
3	60	1.7	4	16.8	4.36	0.18	40.8	45.7
4	74	3.7	4	8.3	2.49	0.20	31.2	31.9
5	61	4.2	4	24.1	1.78	0.05	122.2	156.9
6	64	2.5	4	8.4	3.38	0.28	26.9	27.8
7	58	2.9	4	12.5	2.62	0.14	54.9	58.5
8	62	3.3	4	13.1	2.00	0.10	69.1	74.8
mean	63.5	2.9	3.5	12.6	2.63	0.16	54.0	60.6
SD	5.0	0.7	0.9	5.5	0.85	0.06	30.5	41.6

TABLE 4. RED BLOOD CELL PHARMACOKINETIC PARAMETERS AFTER A SINGLE ORAL DOSE OF 500MG QUININE BASE WITH 4MG CHLORPHENIRAMINE.

Subject	Weight kg	Cmax mg L-1	tmax h	t _{1/2} hL-1	AUC _{0-t} mgh	AUC _{0-t} mghL-1	AUCRBC _{0-t} AUC Plasma _{0-t}
1	67	1.4	4	8.8	9.68	12.23	0.14
2	62	1.0	4	15.1	16.25	20.62	0.38
3	60	1.9	12	-	15.73	15.73	0.29
4	74	1.7	3	18.5	23.29	28.62	0.90
5	61	1.1	8	17.7	26.10	31.20	0.23
6	64	1.4	1	15.4	15.40	19.74	0.40
7	58	1.4	10	18.7	31.39	39.48	0.43
8	62	1.0	6	19.7	20.68	26.36	0.25
mean	63.5	1.35	6	16.27	19.81	24.24	0.38
SD	5.0	0.32	3.7	3.71	6.94	8.89	0.23

TABLE 5a. PLASMA PHARMACOKINETIC PARAMETERS AFTER A SINGLE ORAL DOSE OF 500MG QUININE BASE WITH 250MG TETRACYCLINE.

Subject	Weight kg	Cmax mg L ⁻¹	tmax h	t _{1/2} hL ⁻¹	Vz L kg ⁻¹	CLP Lhkg ⁻¹	AUC _{0-t} mgh L ⁻¹	AUC _{0-∞} mghL ⁻¹
1	67	2.6	6.0	15.8	2.47	0.10	58.6	67.7
2	62	5.6	2.0	9.1	1.93	0.14	52.5	53.8
3	60	2.2	3.0	25.7	3.72	0.10	59.6	81.9
4	74	1.1	6.0	13.4	5.04	0.26	21.5	25.3
5	61	3.7	6.0	22.3	2.23	0.06	90.5	116.3
6	64	1.9	4.0	9.9	2.85	0.19	31.9	38.5
7	58	2.6	3.0	12.8	2.94	0.16	49.3	53.0
mean	63.7	2.8	4.3	15.5	3.02	0.14	51.9	62.3
SD	5.3	1.4	1.7	6.2	1.00	0.06	22.0	30.0

TABLE 5b. PLASMA PHARMACOKINETIC PARAMETERS AFTER A SINGLE ORAL DOSE OF 500MG QUININE BASE ALONE.

Subject	Weight kg	C _{max} mg L ⁻¹	t _{max} h	t _{1/2} hL ⁻¹	V _z L kg ⁻¹	CLP Lhkg ⁻¹	AUC _{0-t} mgh L ⁻¹	AUC _{0-∞} mghL ⁻¹
1	67	2.8	2.0	7.3	1.90	0.18	39.3	40.3
2	62	3.0	2.0	10.2	2.39	0.16	47.4	48.9
3	60	1.7	4.0	16.6	4.29	0.18	41.0	45.8
4	74	3.0	4.0	7.6	2.33	0.21	30.2	31.3
5	61	4.2	4.0	24.1	1.78	0.05	122.2	56.9
6	64	2.3	4.0	9.3	3.71	0.27	25.5	26.7
7	58	2.9	4.0	12.5	2.61	0.14	54.9	58.6
mean	63.7	2.8	3.4	12.5	2.71	0.17	51.5	58.3
SD	5.3	0.7	0.9	6.0	0.93	0.06	52.6	44.7